arXiv:1207.1441v1 [cond-mat.quant-gas] 5 Jul 2012

Thermally isolated Luttinger liquids with noisy Hamiltonians

Luca D'Alessio[1] and Armin Rahmani[2]
[1] *Physics Department, Boston University, Boston, MA 02215, USA*
[2] *Theoretical Division, T-4 and CNLS, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

(Dated: July 9, 2012)

We study the dynamics of a quantum-coherent thermally isolated Luttinger liquid with noisy Luttinger parameter. We calculate the out-of-equilibrium energy fluctuations as well as the correlation functions of this noise-driven system, and find universal features. By identifying two types of energy moments, which can help tease apart the effects of classical and quantum sources of fluctuations, we argue that the classical source is dominant in the thermodynamic limit.

PACS numbers: 05.40.-a, 05.70.Ln, 37.10.Jk, 71.10.Pm

Recent experimental developments with ultracold atoms have motivated numerous studies of the nonequilibrium dynamics of thermally isolated many-body quantum systems. Most of these studies focus on *deterministic* quantum evolution, generated, e.g., by a sudden quench or gradual ramping of the Hamiltonian (see Ref. [1] and the references therein). However, *stochastic* driving of thermally isolated systems with noisy Hamiltonians [2–4] is much less studied, and, in particular, the role of quantum coherence remains largely unexplored. As shown in Fig. 1, fluctuations in such systems stem from two distinct sources: i) the (classical) stochastic nature of the driving (different realizations of noise result in different wave functions with different energy expectation values) and ii) inherent quantum fluctuations (each wave function can be a coherent superposition of states with different energies).

The general setup of the problem is as follow. We consider a quantum system with Hamiltonian $H_0 = \sum_i g_i \hat{O}_i$, where g_is represent the coupling constants and $\hat{O}_i$s are some local operators. The system is initially at zero temperature, i.e., in the ground state of H_0, where the initial energy distribution (see Fig. 1) is a delta function. For $t > 0$, the coupling constants fluctuate in time: $g_i(t) = g_i + \delta g_i(t)$, where, by assumption, $|\delta g_i(t)| \ll |g_i|$. For each realization $\delta g_i(t)$ of noise, the system is then described by a pure state, i.e., a coherent superposition, which evolves unitarily with Hamiltonian $H(t) = H_0 + \sum_i \delta g_i(t)\hat{O}_i$.

Such noisy systems can be experimentally realized by cold atoms in optical potentials, with $\delta g_i(t)$ generated by small fluctuations in the potential. For each realization of noise, the atomic system *sees* an effective time-dependent Hamiltonian $H(t)$, which does not generically commute with H_0. Thus, the quantum evolution creates excitations with respect to the ground state of H_0, increasing the system's energy. Note that the driving with $H(t)$ is assumed external, i.e., there is no feedback action from the quantum system on $\delta g_i(t)$. Therefore, in isolation from a thermal environment (without a mechanism for dissipation), the system can absorb energy *ad infinitum.*



FIG. 1: (color online). The time evolution of $\langle\epsilon\rangle$ for two different realizations, I and II, of noise. Each $\langle\epsilon\rangle$ is the average of a quantum energy distribution. At $t = 0$, the system is in the ground state for all realizations of noise. a) Neither of the classical or quantum fluctuations dominate. b) Classical fluctuations dominate.

The absorbed energy of such noise-driven thermally isolated systems is a *time-dependent* random variable ϵ (as opposed to the steady states which emerge in noise-driven systems coupled to a heat bath [5]). What is the precise meaning of ϵ for systems described by coherent superpositions of energy eigenstates? How can we characterize the average $\mathrm{E}(\epsilon)$ and the variance $\mathrm{Var}(\epsilon)$ of this random variable as a function of time? How are the correlation functions affected by this stochastic driving?

Our objective in this paper is to address the above questions. To do so, we first identify two different types of quantum- and noise-averaged moments, which make the notion of a random absorbed energy precise, and, help tease apart the effects of the classical and quantum sources of fluctuations discussed above (see Fig. 1). We then perform explicit analytical calculations of these moments, as well as different noise-averaged correlation functions, for generic one-dimensional systems described by the Luttinger-liquid (LL) theory.

Our main results are as follows. We find that, in the thermodynamic limit, the classical source dominates the energy fluctuations. In this limit, and in the regime of validity of the LL description, we find a universal relationship (independent of the strength of noise and the Luttinger parameter) between the average and the variance of energy:

$$\mathrm{Var}(\epsilon) = \mathcal{F}(\pi t) \left[\mathrm{E}(\epsilon)\right]^2 , \tag{1}$$

where $\mathcal{F}(x)$ is a dimensionless function of $x = \pi t u/a$ (with velocity u and lattice spacing a set to unity), which decays as $1/x$ for large x. We also find that the noise-induced correction to equal-time correlation functions of scaling dimension Δ, $\mathcal{C}^{(0)}(x) \sim x^{-\Delta}$, has the following universal form:

$$\mathcal{C}(x) - \mathcal{C}^{(0)}(x) \propto W^2 t^3 x^{-\Delta-4}, \tag{2}$$

where W^2 is a constant with dimension of time [see Eq. (8) below], which characterizes the strength of noise.

Let us now discuss the two types of aforementioned energy moments. As the evolution with different noise realizations can lead to different pure-state wave functions, it is helpful to first introduce the notion of a wave-function probability distribution $f(\psi, t)$. We can simply think of $f(\psi, t)$ as the joint probability distribution for all the independent real numbers required to label $|\psi\rangle$ modulo an overall $U(1)$ phase. Similarly, $\int d\psi$ is shorthand for integration over all such numbers. In the scenario considered here (the system initially in the ground state), $f(\psi, t)$ is a delta function at $t = 0$, which broadens for $t > 0$. Note that the wave functions ψ are, generically, coherent superpositions of all energy eigenstates: $|\psi\rangle = \sum_n c_n^\psi |n\rangle$, where $H_0|n\rangle = \epsilon_n|n\rangle$.

The fluctuations of measured energies (with respect H_0) are then characterized by the following moments: $\overline{\langle \epsilon^m \rangle} = \sum_n \epsilon_n^m \mathcal{P}(\epsilon_n)$, where the brackets (overline) indicate a quantum (noise) average, and $\mathcal{P}(\epsilon_n) = \int d\psi f(\psi, t) |c_n^\psi|^2$ is the probability of measuring ϵ_n. This yields

$$\overline{\langle \epsilon^m \rangle} = \int d\psi f(\psi, t) \langle \psi | H_0^m | \psi \rangle = \mathrm{tr}\left[H_0^m \rho(t)\right], \tag{3}$$

where $\rho(t) \equiv \int d\psi f(\psi, t) |\psi\rangle\langle\psi|$ is the density matrix at time t. The fluctuations described by Eq. (3) mix both classical and quantum contributions, which is also the case in experiments.

However, we can alternatively consider the fluctuations of the expectation value of energy, $\langle \epsilon \rangle \equiv \langle \psi | H_0 | \psi \rangle$, over different realizations of noise, which are characterized by

$$\overline{\langle \epsilon \rangle^m} = \int d\psi f(\psi, t) \left(\langle \psi | H_0 | \psi \rangle\right)^m. \tag{4}$$

As $\langle \epsilon \rangle$ is a quantum-averaged quantity, its fluctuations stem solely from the classical stochastic driving. Thus, the difference between these two types of moments can serve as a theoretical diagnostic for the relative importance of the two sources of fluctuations (see Fig. 1). Note that the moments above [Eq. (4)] can not be written in terms of the density matrix for $m > 1$ [6, 7]. Also notice that fluctuations of quantities other than energy can be similarly studied by expanding the wave function in the corresponding eigenbasis.

Let us now turn to the the specific model studied in this paper, namely, a Luttinger Liquid, which is a universal low-energy description of interacting fermions and bosons in one dimension. Since a noise-driven dissipationless system can keep absorbing energy, this low-energy description will eventually break down for most experimentally realistic scenarios. In this paper, we focus on dynamics over a finite time scale where the LL description remains valid. Notice that such time scales can be extended by decreasing the strength of noise. Luttinger liquids with negligible coupling to the environment, which sustain quantum coherence in their life time, have been experimentally realized with cold atoms [8–11].

In terms the Luttinger parameter K and velocity u, the LL Hamiltonian is given by

$$H(K) = u \sum_{q>0} \left(K\, \Pi_q \Pi_{-q} + \frac{1}{K}\, q^2\, \Phi_q \Phi_{-q} \right), \tag{5}$$

where Φ_q are bosonic fields and Π_q their conjugate momenta. The Hamiltonian above can be written as $\sum_{q>0} \left(H_q^{\Re} + H_q^{\Im}\right)$, where $H_q^{\Re\,(\Im)}$ is the Hamiltonian of a single harmonic oscillator involving only the real (imaginary) part of Φ_q. It is convenient to shift the Hamiltonian by a constant, i.e., $H_q^{\Re\,(\Im)} \to H_q^{\Re\,(\Im)} - uq/2$, so that the energies are measured with respect to the ground state. We consider a system initially in the ground state of $H_0 = H(K_0)$, which evolves with $H(K(t)) = H(K_0 + \delta K(t))$ for $t > 0$, where $\delta K(t) \ll K_0$ represents the noise. As the fluctuations of velocity u correspond to a trivial rescaling of the Hamiltonian, we set u to unity throughout this paper.

Expanding the Hamiltonian (5) in δK results in quadratic (in the bosonic fields) noise terms. Thus, integrating out the noise at the outset (as in Refs. [3, 5, 12]) leads to an interacting (quartic) effective action, which is difficult to treat exactly. We thus take an alternative approach described below. As the time-dependent Luttinger parameter is assumed spatially uniform, momentum is a good quantum number, and, the ground state of H_0 is a direct product of Gaussian wave functions. Since each mode evolves with a quadratic Hamiltonian, the time-dependent wave function of the system, i.e., the solution of the time-dependent Schrödinger equation, retains the form

$$\Psi(\{\Phi_q\}, t) = \prod_q \left(\frac{2\, q\, [\Re\, z_q(t)]}{\pi} \right)^{\frac{1}{2}} \exp\left(-q\, z_q(t)\, |\Phi_q|^2\right), \tag{6}$$

where the parameter $z_q(t)$ now satisfies the Riccati equation $i\dot{z}_q(t) = \frac{q}{K(t)} \left[(K(t)\, z_q(t))^2 - 1\right]$, with initial condition $z_q(0) = K_0^{-1}$ [13, 14]. Note that each single-mode wave function in Eq. (6) is exponentially localized on low-energy levels, which acts as an effective high-energy cutoff [15].

Although different modes are decoupled in the Riccati equations above, different $z_q(t)$ evolve with the *same* $K(t)$. Therefore, noise-averaged many-mode quantities must be computed by taking the noise-induced correlations into account. By expanding the above Riccati equation in δK, we then obtain the following nonlinear Langevin equation:

$$i\dot{z}_q = \frac{q}{K_0}\left(K_0^2 z_q^2 - 1\right) - q\left(K_0^2 z_q^2 + 1\right)\delta\alpha, \tag{7}$$

where $\delta\alpha(t) = -\delta K(t)/K_0^2$. Note that there is no dissipative term in the above equation.

We now assume Gaussian noise with zero average and second moments characterized by strength W and correlation time τ as in the Ornstein-Uhlenbeck process:

$$\overline{\delta\alpha(t_1)\delta\alpha(t_2)} = \frac{W^2}{2\tau} e^{-|t_1-t_2|/\tau}. \tag{8}$$

In the limit $\tau \to 0^+$, the right-hand side of Eq. (8) reduced to $W^2\delta(t_1 - t_2)$ describing Gaussian white noise. We leave a discussion of the effects of colored noise (finite τ) to the supplemental material [15], and only consider the $\tau \to 0^+$ limit here. As the white noise is thought of as a limit of a continuous process, we use the Stratonovich interpretation for Eq. (7) [16].

The stochastic differential equation described by Eqs. (7) and (8) is one of the key equations of this paper: it governs the stochastic evolution of $z_q(t)$, which in turn determines the many-body wave function [Eq. (6)], and, consequently, all the observables of the system. We can write observables such as $\langle H_q^m(K)\rangle$, where we have suppressed the $\Re$ and $\Im$ superscripts, in terms of z_q by simple integration over the bosonic fields Φ_q. Similarly, equal-time correlation functions can be expressed in terms of z_q [15]. For example:

$$\langle\Phi(x)\Phi(x')\rangle = \frac{1}{L}\sum_{q>0}\frac{\cos\left[q(x-x')\right]}{q\Re z_q}. \tag{9}$$

As we are interested in the low-energy limit of $|\delta z_q(t)| \ll K_0^{-1}$, where $z_q(t) = K_0^{-1} + \delta z_q(t)$, we can linearize Eq. (7) in δz_q and obtain a simple linear equation [13]:

$$i\,\delta\dot{z}_q = 2q\,\left(\delta z_q - \delta\alpha\right), \tag{10}$$

which admits the explicit solution $\delta z_q(t) = 2iq\int_0^t dt'\ e^{2iq(t'-t)}\delta\alpha(t')$ (we will present an exact treatment of the nonlinear equation later). Our strategy for computing the noise average of generic functions of z_q is then as follows. We expand these functions to leading order in δz_q, and insert the explicit solution above (in terms of $\delta\alpha$) into the resulting expressions. The averaging over noise can then be done by using the Wick's theorem, which relates $\overline{\delta\alpha(t_1)\delta\alpha(t_2)\ldots\delta\alpha(t_n)}$ to two-point functions (8). (As the n-point function above vanishes for odd n, we may need to expand our function to one higher order in δz_q.) Finally, we perform the required integrations over the time arguments to find the noise average.

Let us now compute the two types of energy moments. Here, we only consider the first and the second moments (see the supplement for a discussion of higher moments [15]). As stated above, all observables, including $\langle H_q^m(K_0)\rangle$, can be written in terms of z_q. The expressions for $\langle H_q^m(K_0)\rangle$ become more and more involved as m increases, but if we expand these expressions to leading order in δz_q, we find the simple relationship $\langle H_q^m(K_0)\rangle = q^m\,2^{m-3}\,K_0^2|\delta z_q|^2 + \mathcal{O}(\delta z_q^3)$. Using the strategy outlined after Eq. (10), we then obtain $\overline{\langle H_q^m\rangle} \approx 2^{m-1}q^{m+2}K_0^2W^2t$ and $\overline{\langle H_q\rangle^2} \approx q^6K_0^4W^4\left[2t^2 + \sin^2(2qt)/4q^2\right]$ [15]. Clearly, for a single mode, the energy fluctuations are significantly affected by the quantum source at least in the limit of small excess energy.

Let us now consider the many-mode LL. The average energy can be written as $\overline{\langle\epsilon\rangle} = 2\sum_{q>0}\overline{\langle H_q\rangle}$, where the factor of 2 accounts for the contributions of $H_q^{\Re}$ and $H_q^{\Im}$. Using $\sum_{q>0} \to \frac{L}{2\pi}\int_0^\pi$, we then obtain

$$\overline{\langle\epsilon\rangle} \approx L\pi^3\,K_0^2W^2t/4. \tag{11}$$

We now relate the two types of second moments as follows. We have $\langle\epsilon^2\rangle \equiv \langle H^2\rangle$ and $\langle\epsilon\rangle^2 \equiv \langle H\rangle^2$, with $H = \sum_{q>0}\left(H_q^{\Re} + H_q^{\Im}\right)$. These two moments can then be written, respectively, as sums over q_1 and q_2 of terms of the form $\langle H_{q_1}H_{q_2}\rangle$ and $\langle H_{q_1}\rangle\langle H_{q_2}\rangle$, which only differ when $q_1 = q_2$ and both H_{q_1} and H_{q_2} are in the same $\Re$ or $\Im$ sector (the wave function is a direct product). This leads to

$$\overline{\langle\epsilon^2\rangle} = \overline{\langle\epsilon\rangle^2} + 2\sum_q\left(\overline{\langle H_q^2\rangle} - \overline{\langle H_q\rangle^2}\right). \tag{12}$$

In the above expression, both $\overline{\langle\epsilon\rangle^2}$ and $\overline{\langle\epsilon^2\rangle}$ scale as L^2, while the sum over q scales as L. Therefore, if we take the thermodynamic limit before any other limit, the two types of moments will be, to leading order, identical. As we will see below, this is also the case for the two types of cumulants obtained by subtracting $\left[\overline{\langle\epsilon\rangle}\right]^2$ from the moments above. Such subtraction does not change the scaling with L^2 as an explicit calculation yields [15]:

$$\overline{\langle\epsilon\rangle^2} - \left[\overline{\langle\epsilon\rangle}\right]^2 = \frac{1}{16}\pi^6K_0^4W^4t^2L^2\mathcal{F}(\pi t), \tag{13}$$

where the function $\mathcal{F}(x)$ has the following asymptotic behavior: $\mathcal{F}(x) \approx 2\left(1 - \frac{4}{9}x^2\right)$ for $x \ll 1$, and $\mathcal{F}(x) \approx \frac{16\pi}{7x}$ for $x \gg 1$. This indicates that the cumulant above crosses over from quadratic growth in t for short times to linear growth in t at longer times. Combining Eqs. (13) and (11) leads to the important relationship (1). Recall that the regime of validity for these results can be extended by decreasing the strength of the noise W.

We now turn to the effects of noise on the correlation functions. We compute the noise average of correlation

Correlation function	Ground state	Correction due to noise
$\overline{\langle\Phi(x)\Phi(x')\rangle}$	$\ln(x-x')$	$\propto W^2t^3(x-x')^{-4}$
$\overline{\langle\partial_x\Phi(x)\partial_{x'}\Phi(x')\rangle}$	$(x-x')^{-2}$	$\propto W^2t^3(x-x')^{-6}$
$\overline{\langle e^{i\nu\Phi(x)}e^{-i\nu\Phi(x')}\rangle}$	$(x-x')^{-\nu^2/2}$	$\propto W^2t^3(x-x')^{-4-\nu^2/2}$

TABLE I: The leading scaling behavior of the noise-induced correction at time t to equal-time correlation functions. Standard ground-state behavior ($t=0$) is also shown.

function (9), as well as those of current, $\partial_x\Phi(x)$, and vertex, $e^{i\nu\Phi(x)}$, operators, which, for one realization of noise, can be derived from Eq. (9) by simple differentiation and exponentiation. We are interested in the asymptotic limit $1 \ll x-x' \ll L$ (in units of the lattice spacing). As the LL description is valid for finite t, we also assume $t \ll x$. Again, we use the method outlined after Eq. (10). The calculations are tedious but straightforward [15], and the results for the scaling behavior of the corrections to these noise-averaged correlation functions are summarized in Table. I. Noting that a logarithm has scaling dimension zero, we find that these corrections exhibit the universal feature, Eq. (2).

An alternative approach, which allows us to go beyond the limit of small δz_q, is through the Fokker-Planck (FP) equation for the wave-function probability distribution. Let us first briefly review the general formalism [17]: for a vector $\vec{a}$ of stochastic variables satisfying the Langevin equation $\partial_t a_i = h_i(\vec{a}) + g_i(\vec{a})\gamma(t)$, where h_i and g_i are arbitrary functions of $\vec{a}$ and $\overline{\gamma(t)\gamma(t')} = 2\delta(t-t')$, the probability distribution $f(\vec{a},t)$ evolves according to the FP equation $\partial_t f = \mathcal{D}f$, with the differential operator $\mathcal{D} = -\frac{\partial}{\partial a_i}h_i - \frac{\partial}{\partial a_i}\frac{\partial g_i}{\partial a_j}g_j + \frac{\partial}{\partial a_i}\frac{\partial}{\partial a_j}g_i g_j$ (summation over repeated indices is implied). The noise-averaged expectation value of an arbitrary function $G(\vec{a})$ of the stochastic variables $\vec{a}$ can then be computed at time t as an integral over $\prod_i d\vec{a}_i$ weighted by the formal solution of the FP equation, $f(\vec{a},t) = e^{\mathcal{D}t}f(\vec{a},0)$.

In analogy with the Heisenberg picture of quantum dynamics, we can evolve the observable $G(\vec{a})$ instead of the distribution function $f(\vec{a})$ (using repeated integration by parts), and write

$$\overline{G(\vec{a})}\big|_t = \int \prod_i d\vec{a}_i\, f(\vec{a},0)\, e^{\mathcal{D}^\dagger t}G(\vec{a}), \tag{14}$$

where $\mathcal{D}^\dagger = \left(h_i + \frac{\partial g_i}{\partial a_j}g_j\right)\frac{\partial}{\partial a_i} + g_i g_j \frac{\partial}{\partial a_i}\frac{\partial}{\partial a_j}$. By expanding the exponential operator above as $e^{\mathcal{D}^\dagger t} = \sum_n \frac{t^n}{n!}{\mathcal{D}^\dagger}^n$, we can then compute the noise average of $G(\vec{a})$ as a Taylor expansion in t. Since we are interested in finite time scales, such expansion is indeed very useful even when truncated at a finite order.

For the average energy $\overline{\langle\epsilon\rangle}$, it turns out, however, that we can resum the Taylor series and obtain an exact solution. As the energy is just a sum of the single-mode energies, we need a FP equation for a single mode, i.e., $\vec{a} = (\Re z_q, \Im z_q)$. We can then explicitly construct the differential operator $\mathcal{D}^\dagger$ corresponding to Eq. (7) [15], and find the nth order term by applying it n times to the exact expression for $\langle H_q\rangle$, which can be easily written in terms of $\vec{a}$ [15]. Note that the integration over a_i in Eq. (14) is trivial due to a delta-function initial distribution, $f(\vec{a},0) = \delta(a_1 - K_0^{-1})\delta(a_2)$. By explicitly computing the terms in such expansion to order $n=10$, we found that the series matches a simple exponential, $\overline{\langle H_q\rangle} = \frac{q}{2}\left[\exp\left(2q^2W^2K_0^2t\right) - 1\right]$, term by term. Modeling Eq. (7) by a sequence of random quenches [13, 18], we have also checked this expression numerically. Upon integration over q, we then obtain

$$\overline{\langle\epsilon\rangle} = \frac{L}{8\pi K_0^2W^2t}\left(e^{2K_0^2\pi^2W^2t} - 2\pi^2K_0^2W^2t - 1\right). \tag{15}$$

As expected, the above expression, up to first order in time, agrees with Eq. (11).

Using the Wigner-function approach [19] (see also [20]), we can additionally show that for a thermal initial state (which is subsequently decoupled from the thermal environment during the evolution), the absorbed single-mode energy above is simply multiplied by a prefactor $\coth(q/2k_BT_0)$, where T_0 is the initial temperature, and k_B the Boltzmann constant [15]. The many-mode energy can then be similarly computed by integration over q. Our FP approach is most useful for quantities such as the first and the second moments of energy and current-current correlation functions, where the noise-induced correlations do not need to be treated simultaneously for a large number of modes. The vertex-operator correlation functions, e.g., are not easily tractable with this method. Note that, to compute the second moment of the total energy, one needs to construct a FP equation for four stochastic variable, $\vec{a} = (\Re z_{q_1}, \Im z_{q_1}, \Re z_{q_2}, \Im z_{q_2})$, and so on and so forth for higher moments.

In summary, we studied a thermally isolated LL, randomly driven with a noisy Luttinger parameter, and undergoing coherent quantum evolution for each realization of noise. We computed noise-averaged correlation functions, and studied the energy fluctuations. We characterized these fluctuations by two types of energy moments: one that mixes the classical and quantum sources of fluctuations and one that is only affected by the classical source. We found that while for a single mode, the two types of moments lead to very different results, for the many-mode problem, the difference disappears in the thermodynamic limit. This indicates that many-body properties of such noise-driven coherent systems likely exhibit effective decoherence. Our approach to the dynamics of noisy LLs is based on a mapping of the quantum problem to one of nonequilibrium classical statistical mechanics, which provides powerful tools, such as the FP equation, for performing exact calculations.

We are grateful to K. Barros, C. Chamon, A. del

Campo, T. Giamarchi, D. Huse, P. Krapivsky, I. Martin, A. Polkovnikov, P. Zoller and W. Zurek for helpful discussions. This work was supported in part by AFOSR FA9550-10-1-0110 (LD) and the U.S. Department of Energy through LANL/LDRD program (AR).

[1] A. Polkovnikov, K. Sengupta, A. Silva and M. Vengalattore, Rev. Mod. Phys. **83**, 863 (2011).
[2] G. Bunin, L. D'Alessio, Y. Kafri and A. Polkovnikov, Nat. Phys. **7**, 913 (2011).
[3] J. Marino and A. Silva, arXiv:1203.2108v2 [cond-mat.stat-mech].
[4] H. Pichler, J. Schachenmayer, J. Simon, P. Zoller and A. J. Daley, arXiv:1205.6189v1 [cond-mat.quant-gas].
[5] E. G. Dalla Torre, E. Demler, T. Giamarchi and E. Altman, Nat. Phys. **6**, 806 (2010); E. G. Dalla Torre, E. Demler, T. Giamarchi and E. Altman, Phys. Rev. B **85**, 184302 (2012).
[6] K. Mølmer, Y. Castin and J. Dalibard, J. Opt. Soc. Am. B **10**, 524 (1993).
[7] H.-P. Breuer and F. Petruccione, The theory of open quantum systems, Oxford University Press, Oxford (2006).
[8] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Flling, I. Cirac, G. V. Shlyapnikov, T. W. Hnsch and I. Bloch, Nature **429**, 277 (2004).
[9] T. Kinoshita, T. Wenger and D. S. Weiss, Science **305**, 1125 (2004).
[10] A. Recati, P. O. Fedichev, W. Zwerger and P. Zoller, Phys. Rev. Lett. **90** 020401 (2003).
[11] A. Recati, P. O. Fedichev, W. Zwerger and P. Zoller, J. Opt. B. **5** S55 (2003).
[12] J. H. Wilson, B. M. Fregoso and V. M. Galitski, Phys. Rev. B **85**, 174304 (2012).
[13] A. Rahmani and C. Chamon, Phys. Rev. Lett. **107**, 016402 (2011).
[14] A. Polkovnikov and V. Gritsev, Nat. Phys. **4**, 477 (2008).
[15] See supplemental material for details.
[16] P. E. Koeden, P. Platen, *Numerical Solution of Stochastic Differential Equations*, Applications of Mathematics, Stochastic Modelling and Applied Probability, 23, Springer-Verlag, Berlin, 1992.
[17] H. Risken, The Fokker-Planck Equation: Methods of Solutions and Applications, Springer (1989).
[18] E. Perfetto, G. Stefanucci and M. Cini, Phys. Rev. Lett. **105**, 156802 (2010); E. Perfetto and G. Stefanucci, Euro. Phys. Lett **95**, 10006 (2011).
[19] E. Wigner, Phys. Rev. **40**, 749 (1932).
[20] A. Polkovnikov, Ann. Phys. **325**, 1790 (2010).

Supplemental Material for “Thermally isolated Luttinger liquids with noisy Hamiltonians”

Luca D’Alessio[1] and Armin Rahmani[2]
[1]*Physics Department, Boston University, Boston, MA 02215, USA*
[2]*Theoretical Division, T-4 and CNLS, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

(Dated: July 9, 2012)

arXiv:1207.1441v1 [cond-mat.quant-gas] 5 Jul 2012

I. EIGENSTATE EXPANSION OF THE SINGLE-MODE WAVE FUNCTION

In this section, we compute the overlaps of the single-mode wave function with the eigenstates of the single-mode Hamiltonian H_q. The wave function is given by

$$\psi_q(\phi,t)=\left(\frac{2\,q\,[\Re\,z_q(t)]}{\pi}\right)^{\frac{1}{4}}\exp\left(-q\,z_q(t)\,\phi^2\right), \tag{1}$$

where ϕ represents either $\Phi_q^{\Re}$ or $\Phi_q^{\Im}$ [see Eq. (6) of the main text]. Analogously, the Hamiltonian H_q is shorthand for either $H_q^{\Re}$ or $H_q^{\Im}$:

$$H_q(K(t))=u\left(-\frac{K(t)}{4}\,\partial_\phi^2+\frac{1}{K(t)}\,q^2\phi^2\right). \tag{2}$$

In the absence of noise, the Hamiltonian is given by $H_0\equiv H_q(K_0)$. For $z_q=1/K_0$, the wave function (1) is the ground state of Hamiltonian (2). For arbitrary z_q, however, it is a superposition $\psi_q(\phi)=\sum_{n=0}^{\infty}c_n(z_q)\psi_n(\phi)$, where $\psi_n(\phi)$ is an eigenfunction of H_0 with energy $\left(n+\frac{1}{2}\right)uq$, which can be written explicitly in terms of the Hermite polynomials. Through direct integration, we can compute the amplitudes $c_n(z_q)=\int d\phi\,\psi_q(\phi)^\star\psi_n(\phi)$, which yield:

$$|c_{2n+1}(z_q)|^2=0,\ \ |c_{2n}(z_q)|^2=\frac{(2n)!}{2^{2n-1}n!n!}\,\frac{x_q^{1/2}\left[(x_q-1)^2+y_q^2\right]^n}{\left[(x_q+1)^2+y_q^2\right]^{n+1/2}},$$

where

$$x_q\equiv K_0\left[\Re\,z_q(t))\right],\qquad y_q\equiv K_0\left[\Im\,z_q(t)\right].$$

We see from the above expression that the overlaps $|c_n(z_q)|$ are identically zero for odd n, and decay exponentially for even n.

II. EXPRESSIONS FOR OBSERVABLES

In this section, we express some observables in terms of $z_q(t)$. Measuring the energies with respect to the ground state of $H_q(K_0)$, and using the explicit form of the wave function (1), we can obtain $\langle H_q(K_0)\rangle$ by direct integration over the bosonic fields:

$$\langle H_q(K_0)\rangle\equiv\int_{-\infty}^{+\infty}d\psi\;\psi_q(\phi,t)^\star H_q(K_0)\psi_q(\phi,t)=\frac{q}{2}\left[\frac{1}{2K_0\Re z_q}\left(1+K_0^2|z_q|^2\right)-1\right], \tag{3}$$

where u is set to unity. Similarly we can compute the second moment of energy:

$$\langle H_q^2(K_0)\rangle=\langle H_q(K_0)\rangle^2+\frac{q^2}{8K^2\Re z_q^2}\left[1-2K_0^2(\Re z_q^2-\Im z_q^2)+K_0^4|z_q|^4\right]. \tag{4}$$

Note that both expressions above vanish for the ground state ($z_q=K_0^{-1}$). This simple approach can be used to compute higher moments of energy, but the resulting expressions become more cumbersome. Using MATHEMATICA®, we have calculated the moments $\langle H_q^m(K_0)\rangle$ for $m\le 20$, and checked that to leading order in $\delta z_q=z_q-K_0^{-1}$, they can be written as:

$$\langle\epsilon_q^m\rangle=\langle H_q^m(K_0)\rangle=q^m\,2^{m-3}\,K_0^2|\delta z_q|^2+\mathcal{O}(\delta z_q^3). \tag{5}$$

Equal-time correlation functions are also simple to compute in term of $z_q(t)$. For example, Eq. (9) of the main text can be derived as follows. We can expand $\Phi(x)$ in Fourier modes, $\Phi(x) = \sum_q \frac{e^{iqx}}{\sqrt{L}}\Phi_q = \frac{2}{\sqrt{L}}\sum_{q>0}\left[\cos(qx)\Re\Phi_q - \sin(qx)\Im\Phi_q\right]$, and make use of the relation $\langle \Re\Phi_{q_1}\Re\Phi_{q_2}\rangle = \langle \Im\Phi_{q_1}\Im\Phi_{q_2}\rangle = \delta_{q_1 q_2}/4q\Re z_q$ to write

$$\langle \Phi(x)\Phi(x')\rangle = \frac{1}{L}\sum_{q>0}\frac{\cos\left[q(x-x')\right]}{q\Re z_q}. \tag{6}$$

For the ground state, $z_q = K_0^{-1}$, Eq. (6) reduces to its usual asymptotic logarithmic form in the thermodynamic limit ($\sum_{q>0} \to \frac{L}{2\pi}\int_\epsilon^\pi dq$):

$$\langle \Phi(x)\Phi(0)\rangle_{GS} = \frac{K_0}{2\pi}\int_\epsilon^\pi dq\frac{\cos(qx)}{q} = \frac{K_0}{2\pi}(\mathrm{Ci}(\pi x) - \mathrm{Ci}(\epsilon x)), \tag{7}$$

where $\epsilon \simeq \frac{2\pi}{L}$ is a short-distance momentum cutoff, and the cosine integral $\mathrm{Ci}(\zeta) \equiv -\int_\zeta^\infty \frac{\cos t}{t}dt$ has the following asymptotic form for small ζ: $\mathrm{Ci}(\zeta) \approx \gamma + \ln\zeta$, where γ is the Euler-Mascheroni constant. In the asymptotic limit, $1 \ll x \ll L$, where the lattice spacing is set to unity, we can neglect $\mathrm{Ci}(\pi x)$, and recover the usual logarithmic correlation function.

III. HIGHER MOMENTS OF ENERGY

Focusing on a single mode for simplicity, we calculate the noise-averaged moments $\overline{\langle\epsilon_q\rangle^m}$. To leading order in δz_q, we can write $\langle\epsilon_q\rangle^m$ as [see Eq. (5) above]

$$\langle\epsilon_q\rangle^m \approx \left(\frac{K_0^2 q}{4}|\delta z_q|^2\right)^m. \tag{8}$$

After substituting the explicit solution, $\delta z_q(t) = 2iq\int_0^t dt'\, e^{2iq(t'-t)}\delta\alpha(t')$, of the linear Langevin equation [see Eq. (10) of the main text], we need to compute

$$\overline{|\delta z_q|^{2m}} = (2q)^{2m}\int_0^t dt_1 e^{2iqt_1}\int_0^t dt_1' e^{-2iqt_1'}\ldots\int_0^t dt_m' e^{-2iqt_m'} \times \overline{\delta\alpha(t_1)\delta\alpha(t_1')\ldots\delta\alpha(t_m)\delta\alpha(t_m')}. \tag{9}$$

For Gaussian white noise, $\overline{\delta\alpha(t)\delta\alpha(t')} = W^2\delta(t-t')$, the above $2m$-point function breaks up into a sum of products of m delta functions. The number of terms in this sum, i.e., the number of different contractions, is $(2m-1)!!$. There are three different types of contractions: (i) $\overline{\delta\alpha(t_i)\delta\alpha(t_j)}$, (ii) $\overline{\delta\alpha(t_i')\delta\alpha(t_j')}$, and (iii) $\overline{\delta\alpha(t_i)\delta\alpha(t_j')}$. Note that we have an equal number of contractions of type (i) and (ii), which are related to each other by complex conjugation. Upon integration, we obtain

$$\begin{aligned}
&\int_0^t dt_i\int_0^t dt_j e^{2iq(t_i+t_j)}\overline{\delta\alpha(t_i)\delta\alpha(t_j)} \times \int_0^t dt_m'\int_0^t dt_n' e^{-2iq(t_m+t_n)}\overline{\delta\alpha(t_m')\delta\alpha(t_n')} = W^4\left(\frac{\sin(2qt)}{2q}\right)^2,\\
&\int_0^t dt_i\int_0^t dt_j' e^{2iq(t_i-t_j')}\overline{\delta\alpha(t_i)\delta\alpha(t_j')} = W^2 t.
\end{aligned} \tag{10}$$

We can then write

$$\overline{\langle\epsilon_q\rangle^m} \approx q^{3m}\,(K_0^2W^2t)^m\sum_{p=0}^{\lfloor m/2\rfloor} S_{m,p}\left(\frac{\sin 2qt}{2qt}\right)^{2p}, \tag{11}$$

where $\lfloor x\rfloor$ indicates the floor value of x and $S_{m,p}$ represents the number of ways for having p contractions of type (i), p contractions of type (ii), and, consequently, $m-2p$ contractions of type (iii). The simple combinatorial argument below gives

$$S_{m,p} = \left[\begin{pmatrix} m \\ 2p\end{pmatrix}\cdot(2p-1)!!\right]^2(m-2p)!.$$

Out of the m choices for $\delta\alpha(t_i)$, we select $2p$, and then make p contractions among them in $(2p-1)!!$ ways. A similar argument applies to $\delta\alpha(t_i')$, which results in the power of 2 for the term in the square brackets. We now have $m-2p$ unpaired $\delta\alpha(t_i)$, and $m-2p$ unpaired $\delta\alpha(t_i')$, which can be paired up in $(m-2p)!$ ways.

IV. DERIVATION OF EQ. (13) OF THE MAIN TEXT

Let us now explicitly compute $\overline{\langle\epsilon\rangle^2}$. Since the energy is the sum of single-modes energies, $\langle\epsilon\rangle = 2\sum_q \langle H_q\rangle$ (the factor of 2 accounts for the contribution of $H_q^{\Re}$ and $H_q^{\Im}$), we can write $\overline{\langle\epsilon\rangle^2} = 4\sum_{q_1,q_2} \overline{\langle H_{q_1}\rangle\langle H_{q_2}\rangle}$. After inserting Eq. (8) for $m=1$, we obtain

$$\overline{\langle\epsilon\rangle^2} \approx \sum_{q_1,q_2} \frac{K_0^4 q_1 q_2}{4} \overline{|\delta z_{q_1}|^2 |\delta z_{q_2}|^2}, \tag{12}$$

which, upon substituting the explicit solution of the linear Langevin equation [see Eq. (10) of the main text], yields

$$\overline{\langle\epsilon\rangle^2} \approx 4K_0^4 \sum_{q_1,q_2} q_1^3 q_2^3 \iiiint_0^t dt_1 dt_2 dt_1' dt_2' \, e^{2iq_1(t_1-t_1')} e^{2iq_2(t_2-t_2')} \overline{\delta\alpha(t_1)\delta\alpha(t_1')\delta\alpha(t_2)\delta\alpha(t_2')}. \tag{13}$$

As we did in in Sec. III, we use the Wick's theorem to break the 4-point function $\overline{\delta\alpha(t_1)\delta\alpha(t_1')\delta\alpha(t_2)\delta\alpha(t_2')}$ into a sum of products of 2-points functions. However, in Sec. III, we only had one mode q, and two types of contractions. Here, on the other hand, we have three different types of contractions. Upon inserting these delta functions and performing the integrals over time, we find

$$\overline{\langle\epsilon\rangle^2} = 4K_0^4 W^4 \sum_{q_1,q_2} q_1^3 q_2^3 \left\{ \frac{\sin^2[(q_1+q_2)t]}{(q_1+q_2)^2} + \frac{\sin^2[(q_1-q_2)t]}{(q_1-q_2)^2} + t^2 \right\}. \tag{14}$$

The term proportional to t^2 is exactly equal to $\left[\overline{\langle\epsilon\rangle}\right]^2$. To obtain Eq. (13) of the main text, we convert the sums to integrals, and define new variables $q_i = \pi\xi_i$ and $x = \pi t$, which lead to Eq. (13) of the main text with

$$\mathcal{F}(x) = 16 \int_0^1 d\xi_1 \int_0^1 d\xi_2 \, \xi_1^3 \xi_2^3 \left\{ \frac{\sin^2[(\xi_1+\xi_2)x]}{[(\xi_1+\xi_2)x]^2} + \frac{\sin^2[(\xi_1-\xi_2)x]}{[(\xi_1-\xi_2)x]^2} \right\}. \tag{15}$$

For $x \ll 1$, the asymptotic behavior is obtained by a simple Taylor expansion in x, which gives $\mathcal{F}(x) \sim 2\left(1-\frac{4}{9}x^2\right)$. For $x \gg 1$, the main contribution comes from $\xi_1 \approx \xi_2$ then we can approximate $\mathcal{F}(x) \sim 16\,\frac{1}{2}\int dQ \int dq \left(\frac{Q+q}{2}\right)^3 \left(\frac{Q-q}{2}\right)^3 \frac{\sin^2(qx)}{(qx)^2}$ where we have defined $Q = \xi_1+\xi_2$ and $q = \xi_1-\xi_2$ and the factor 1/2 comes from the Jacobian of the transformation. Since $Q \gg q$ the dominant contribution comes from the term proportional to Q^6. The final change of variable $k = qx$ leads to $\mathcal{F}(x) \sim \frac{16}{2^7}\int_0^2 dQ \int_{-\infty}^{+\infty} \frac{dk}{x} \, \frac{Q^6 \sin^2(k)}{k^2} = \frac{16\pi}{7x}$.

V. EFFECTS OF COLORED NOISE

In this section, we discuss the effects of finite correlation time τ in the Ornstein-Uhlenbeck process:

$$\overline{\delta\alpha(t_1)\delta\alpha(t_1')} = \frac{W^2}{2\tau} e^{-|t_1-t_1'|/\tau}. \tag{16}$$

For the single-mode energy moments $\overline{\langle\epsilon_q^m\rangle} \approx q^m 2^{m-3} K_0^2 \, \overline{|\delta z_q|^2}$, we need to evaluate

$$\overline{|\delta z_q|^2} = (2q)^2 \int_0^t dt_1 \int_0^t dt_1' \, e^{2iq(t_1-t_1')} \, \overline{\delta\alpha(t_1)\delta\alpha(t_1')}. \tag{17}$$

Substituting Eq. (16), and performing the integral yields:

$$\overline{(|\delta z_q|^2)} = 4q^2 W^2 t \left[\frac{1}{1+(2q\tau)^2} + \frac{\tau}{t} \Re\left(\frac{e^{-\frac{t}{\tau}(1+2q\tau)} - 1}{(1+2iq\tau)^2} \right) \right]. \tag{18}$$

Note that, for $t \gg \tau$, the above expression simplifies to $\overline{|\delta z_q|^2} = 4q^2 \frac{W^2}{1+(2q\tau)^2}\, t$, while for white noise, $\tau \to 0^+$, we obtain $\overline{|\delta z_q|^2} = 4q^2 W^2 t$. We then conclude that, for $t \gg \tau$, the single-mode energy can be obtained from the corresponding expression for white noise by the simple rescaling $W^2 \to \frac{W^2}{1+(2q\tau)^2}$. The average energy can then be calculated by integration over momentum:

$$\overline{\langle\epsilon\rangle} = 2\sum_q \overline{\langle\epsilon_q\rangle} \approx K_0^2 W^2 t \frac{L}{\pi}\int_0^\pi dq \frac{q^3}{1+(2q\tau)^2} = \frac{L}{\pi}\,\frac{K_0^2 W^2 t}{32\tau^4}\,\left[(2\pi\tau)^2 - \log(1+(2\pi\tau)^2)\right], \tag{19}$$

which for white noise, $\tau \to 0^+$, reduces to $\overline{\langle\epsilon\rangle} = L\pi^3 K_0^2 W^2 t/4$.

Let us now consider the higher moments of the single-mode energy, which are given by $\langle\epsilon_q\rangle^m \approx \left(\frac{K_0^2 q}{4}|\delta z_q|^2\right)^m$ [see Eq. (8)]. Using the method of Sec. III, we arrive at the analogue of Eq. (10), which reads

$$\begin{aligned}
&\int_0^t dt_i \int_0^t dt_j e^{2iq(t_i+t_j)}\overline{\delta\alpha(t_i)\delta\alpha(t_j)} \times \int_0^t dt'_m \int_0^t dt'_n e^{-2iq(t_m+t_n)}\overline{\delta\alpha(t'_m)\delta\alpha(t'_n)} \\
&= W^4\, e^{-2t/\tau}\left[\frac{2q\tau + e^{t/\tau}\left(\sin(2qt) - 2q\tau\cos(2qt)\right)}{2q(1+(2q\tau)^2)}\right]^2, \\
&\int_0^t dt_i \int_0^t dt'_j e^{2iq(t_i-t'_j)}\overline{\delta\alpha(t_i)\delta\alpha(t'_j)} = W^2 t\left[\frac{1}{1+(2q\tau)^2} + \frac{\tau}{t}\Re\left(\frac{e^{-\frac{t}{\tau}(1+2q\tau)}-1}{(1+2iq\tau)^2}\right)\right].
\end{aligned} \tag{20}$$

Note that, for $t \gg \tau$, the two expressions above respectively simplify to

$$\left(\frac{W^2}{1+(2q\tau)^2}\right)^2\left(\frac{\sin(2qt) - 2q\tau\cos(2qt)}{2q}\right)^2, \qquad \frac{W^2}{1+(2q\tau)^2}\,t, \tag{21}$$

while for white noise, we obtain

$$W^4\left(\frac{\sin(2qt)}{2q}\right)^2, \qquad W^2\,t \tag{22}$$

by replacing $\tau = 0$ in the above expressions.

We observe that finite correlation time does not qualitatively change the white-noise behavior. We can now write the analogue of Eq. (11) as

$$\overline{\langle\epsilon_q\rangle^m} \approx \left(q^3 K_0^2 \frac{W^2}{1+(2q\tau)^2}t\right)^m \sum_{p=0}^{\lfloor m/2\rfloor} S_{m,p}\left(\frac{\sin(2qt) - 2q\tau\cos(2qt)}{2qt}\right)^{2p}, \tag{23}$$

where $S_{m,p}$ is defined in Sec. III.

Finally, we generalize Eq. (15). By repeating the derivation in Sec. IV, we arrive at the analogue of Eq. (14), which in the limit of $t \gg \tau$ reduces to

$$\begin{aligned}
\overline{\langle\epsilon\rangle^2} = 4K_0^4 W^4 \sum_{q_1,q_2} q_1^3 q_2^3 \times \Bigg\{ &\frac{\left[t(1+m_1^2) + \tau(m_1^2-1)\right]\left[t(1+m_2^2) + \tau(m_2^2-1)\right]}{(1+m_1^2)^2(1+m_2^2)^2} + \\
&\left(\frac{(q_1+q_2)\tau(1+m_1m_2)\cos\left[(q_1+q_2)t\right] - \left(1+\frac{1}{2}(m_1^2+m_2^2)\right)\sin\left[(q_1+q_2)t\right]}{(q_1+q_2)(1+m_1^2)(1+m_2^2)}\right)^2 + \\
&\left(\frac{(q_1-q_2)\tau(1+m_1m_2)\cos\left[(q_1-q_2)t\right] - \left(1+\frac{1}{2}(m_1^2+m_2^2)\right)\sin\left[(q_1-q_2)t\right]}{(q_1-q_2)(1+m_1^2)(1+m_2^2)}\right)^2 \Bigg\},
\end{aligned} \tag{24}$$

where we have introduced $m_i = 2q_i\tau$. The first term in the curly brackets is equal to $\left[\overline{\langle\epsilon\rangle}\right]^2$. Moreover, it is easy to see that in the limit $\tau \to 0^+$ we recover Eq. (14). Replacing summation over q_i by $\frac{L}{2\pi}\int dq_i$, and making a change of variables:

$$q_i = \pi\xi_i, \qquad x = \pi t, \qquad y = \pi\tau, \tag{25}$$

we obtain

$$\overline{\langle\epsilon\rangle^2} = \left[\overline{\langle\epsilon\rangle}\right]^2 + \frac{1}{16}L^2 K_0^4 W^4 t^2 \pi^6 \mathcal{F}(x,y), \tag{26}$$



FIG. 1: Eq. (27) for $y = \pi\tau = 0,\ 0.1,\ 0.2,\ 0.5$. Note that the equation is valid for $x \gg y$. The $y = 0$ line corresponds to the white-noise case. The asymptotic behavior for $x \ll 1$ and $y = 0$ is also shown (solid black line).

where

$$\begin{aligned}
\mathcal{F}(x,y) \equiv 16 \Bigg\{ \int_0^1 d\xi_1 \int_0^1 d\xi_2 \ \xi_1^3 \xi_2^3 \times \\
\Bigg[\left(\frac{(\xi_1+\xi_2)y(1+4\xi_1\xi_2 y^2)\cos\left[(\xi_1+\xi_2)x\right]-(1+2(\xi_1^2y^2+\xi_2^2y^2))\sin\left[(\xi_1+\xi_2)x\right]}{x(\xi_1+\xi_2)(1+(2\xi_1 y)^2)(1+(2\xi_2 y)^2)}\right)^2 + \\
\left(\frac{(\xi_1-\xi_2)y(1-4\xi_1\xi_2 y^2)\cos\left[(\xi_1-\xi_2)x\right]-(1+2(\xi_1^2y^2+\xi_2^2y^2))\sin\left[(\xi_1-\xi_2)x\right]}{x(\xi_1-\xi_2)(1+(2\xi_1 y)^2)(1+(2\xi_2 y)^2)}\right)^2\Bigg]\Bigg\}.
\end{aligned} \tag{27}$$

Note that this expression is valid only for $t \gg \tau$, which implies $x \gg y$. The asymptotic behavior at long times, $x \gg 1$, can be obtained as for Eq. (15): $\mathcal{F}(x,y) \simeq \frac{16\pi}{7x}\left(1-\frac{56}{9}y^2+\mathcal{O}(y^4)\right)$. This implies that the finite correlation time, $y > 0$, does not change the form of the time dependence and simply lowers, by an overall coefficient, the white-noise ($y = 0$) asymptotic value. The short-time behavior can be seen in Fig. 1 for several values of y. In the physically relevant region, $x \gg y$, the curves show a rapid decay followed by a slower one. This behavior is qualitatively similar to the white-noise case [Eq. (15)], which is also plotted in Fig. 1 for comparison.

VI. NONLINEAR FOKKER-PLANCK EQUATION

In this section, we explicitly derive the differential operator $\mathcal{D}^\dagger$ used in the derivation of Eq. (15) of the main text [see also Eq. (14) of the main text]. Let us define

$$\mathscr{R}_i \equiv \Re z_{q_i}, \qquad \mathscr{I}_i \equiv \Im z_{q_i}.$$

We can then write the nonlinear Langevin equation [Eq. (7) of the main text] as

$$\begin{aligned}
\dot{\mathscr{R}}_i &= 2K_0 q_i\, \mathscr{R}_i \mathscr{I}_i - 2K_0^2 q_i\, \mathscr{R}_i \mathscr{I}_i\, \delta\alpha, \\
\dot{\mathscr{I}}_i &= K_0 q_i\, \left(\mathscr{I}_i^2 - \mathscr{R}_i^2 + K_0^{-2}\right) - K_0^2 q_i\, \left(\mathscr{I}_i^2 - \mathscr{R}_i^2 - K_0^{-2}\right)\, \delta\alpha.
\end{aligned} \tag{28}$$

As the total energy is a sum of single-mode energies, we only need a single-mode Fokker-Planck equation, and we can drop the subscript i. Rescaling $\delta\alpha \to \frac{W}{\sqrt{2}}\gamma(t)$, with $\langle\gamma(t)\gamma(t')\rangle = 2\delta(t-t')$, we can read off f_i and g_i for $i = 1, 2$ (in the notation of the main text):

$$f_1 = 2K_0 q a_1 a_2, \quad g_1 = -\sqrt{2} W K_0^2 q a_1 a_2, \quad f_2 = K_0 q\left(a_2^2 - a_1^2 + K_0^{-2}\right), \quad g_2 = -\frac{W}{\sqrt{2}} K_0^2 q\left(a_2^2 - a_1^2 - K_0^{-2}\right), \tag{29}$$

where $\vec{a} \equiv (a_1, a_2) = (\mathscr{R}, \mathscr{I})$.

In terms of the above f_i and g_i, we can write

$$\mathcal{D}^\dagger = D_1^{(1)}\partial_{a_1} + D_2^{(1)}\partial_{a_2} + D_{1,1}^{(2)}\partial_{a_1}^2 + D_{2,2}^{(2)}\partial_{a_2}^2 + 2D_{1,2}^{(2)}\partial_{a_1}\partial_{a_2}, \tag{30}$$

where

$$D_1^{(1)} = f_1 + g_1\frac{\partial g_1}{\partial a_1} + g_2\frac{\partial g_1}{\partial a_2}, \qquad D_2^{(1)} = f_2 + g_1\frac{\partial g_2}{\partial a_1} + g_2\frac{\partial g_2}{\partial a_2}, \qquad D_{i,j}^{(2)} = g_i g_j. \tag{31}$$

We now consider the single-mode energy [Eq. (3)]. The general expression given in the main text [see Eq. (12)] gives the Taylor series of the noise-averaged quantity: $\overline{\langle \epsilon_q \rangle} = \sum_{n=0} \frac{t^n}{n!} c_n$, where the coefficient c_n is obtained by applying the operator ${\mathcal{D}^\dagger}^n$ to Eq. (3), and setting $\Re z = K_0^{-1}$ and $\Im z = 0$ in the resulting expression. We have explicitly calculated such terms for $n \leq 10$, and found that $c_0 = 0$, and, for $n \geq 1$, $c_n = \frac{qR}{2}\left(2q^2W^2K_0^2\right)^n$. We are thus lead to conjecture that the behavior holds for all n, and the series can be resummed to the exponential expression in the main text. We have also checked this conjecture by numerical simulations. Note that if we are interested in 2-modes quantities, we need to repeat the procedure above with $\vec{a} = (\mathscr{R}_1, \mathscr{I}_1, \mathscr{R}_2, \mathscr{I}_2)$, identify f_i, g_i for $i = 1, 2, 3, 4$, and build the 4-dimensional vector $D^{(1)}$ and the 4×4 matrix $D^{(2)}$. In general, if we need the correlations of m momenta, $D^{(1)}$ is $2m$-dimensional and $D^{(2)}$ is a $2m \times 2m$ matrix.

VII. WIGNER-FUNCTION APPROACH

An alternative approach to the dynamics of the problem is through the Wigner-function representation. This approach allows us to treat, on the same footing, initial conditions given by a finite-temperature thermal density matrix as well as the zero-temperature ground state considered so far. Note that the system is still thermally isolated during the evolution, i.e., it is decoupled from a finite-temperature heat bath at time $t = 0$.

Let us briefly review the formalism. In terms of phase-space variable x and p (which can be vectors for multi-dimensional problems), the Weyl symbol $\Omega_w(x,p)$ for a quantum operator $\hat{\Omega}$ is defined as

$$\Omega_w(x,p) = \int ds \left\langle x - \frac{s}{2}\middle| \hat{\Omega} \middle| x + \frac{s}{2}\right\rangle \exp\left[\frac{i}{\hbar} p \cdot s\right]. \tag{32}$$

The Wigner function $W(x,p)$ is, by definition, the Weyl symbol for the density matrix $\hat{\rho}$, which satisfies the following equation of motion:

$$\partial_t W = -\frac{2}{\hbar} H_w \sin\left(\frac{\hbar}{2}\Lambda\right) W, \tag{33}$$

where H_w is the Weyl symbol for the Hamiltonian, and $\Lambda \equiv \overleftarrow{\partial_p}\,\overrightarrow{\partial_x} - \overleftarrow{\partial_x}\,\overrightarrow{\partial_p}$. The expectation value of an operator $\hat{O}$ can be written as

$$\langle \hat{O} \rangle = \iint_{-\infty}^{+\infty} \frac{dxdp}{2\pi}\, W(x,p)\; O_w(x,p) \tag{34}$$

in terms of its Weyl symbol $O_w(x,p)$ and the Wigner function $W(x,p)$.

Let us now consider the single-mode Hamiltonian $\hat{H}_q(K(t)) = \left(\frac{K(t)}{4}\,\hat{p}^2 + \frac{1}{K(t)}\,q^2\hat{x}^2\right)$ [see Eq. (2)], where we have set $u = 1$ and suppressed the $\Re$ or $\Im$ superscripts. Here, $\hat{x}$ and $\hat{p}$ respectively represent the real (or imaginary) part of operators Φ_q and Π_q, and we have used the hat notation to tell apart quantum operators from phase-space variables. For a system evolving from an initial thermal density matrix,

$$\hat{\rho}_0 = \exp\left(-\hat{H}_q(K_0)/k_B T_0\right)/\mathrm{tr}\left[\exp\left(-\hat{H}_q(K_0)/k_B T_0\right)\right], \tag{35}$$

with the Hamiltonian above (for arbitrary $K(t)$), the Wigner function retains the following form:

$$W(x,p) = \mathcal{N}\exp\left[-\frac{A}{2}x^2 - \frac{B}{2}p^2 + Cxp\right], \tag{36}$$

where A, B, C and $\mathcal{N}$ are potentially time-dependent functions with the following initial conditions:

$$A(t=0) = \frac{4q}{K_0 R}, \quad B(t=0) = \frac{K_0}{qR}, \quad C(t=0) = 0, \quad \mathcal{N}(t=0) = \frac{2}{R}, \quad R = \coth\left(\frac{q}{2k_B T_0}\right). \tag{37}$$

Note that the normalization condition $\iint_{-\infty}^{+\infty} \frac{dxdp}{2\pi} W(x,p) = 1$ sets

$$\mathcal{N} = \sqrt{AB - C^2} \tag{38}$$

at all times. For the quadratic hamiltonian $\hat{H}_q(K(t))$, the Weyl symbol is simply obtained by replacing the operators $\hat{x}, \hat{p}$ with the phase-space variable x, p. We can then write Eq. (33) as

$$W(x,p)\left(\frac{\dot{\mathcal{N}}}{\mathcal{N}} - \frac{\dot{A}}{2}x^2 - \frac{\dot{B}}{2}p^2 + \dot{C}xp\right) = W(x,p)\left(2\frac{q^2}{K(t)}\,x\,(Cx - Bp) - \frac{K(t)}{2}\,p\,(Cp - Ax)\right),$$

which leads to the following equations for the parameters of the Wigner function:

$$\dot{\mathcal{N}} = 0, \quad \dot{A} = -4\frac{q^2}{K(t)}C, \quad \dot{B} = K(t)C, \quad \dot{C} = 2\left(\frac{K(t)}{4}A - \frac{q^2}{K(t)}B\right). \tag{39}$$

By using Eq. (38), and noting that $\mathcal{N}$ does not change from its initial value, we can eliminate A from the equations above:

$$\dot{B} = K(t)C, \qquad \dot{C} = K(t)\frac{(4 + C^2R^2)}{2R^2B} - \frac{2q^2B}{K(t)}. \tag{40}$$

We now expand Eq. (40) in $\delta K(t) \ll K_0$, and obtain two coupled Langevin equations:

$$\begin{aligned} \dot{B} &= CK_0 - CK_0^2\delta\alpha(t), \\ \dot{C} &= \frac{K0(4 + C^2R^2)}{2R^2B} - \frac{2q^2B}{K_0} - \left[2Bq^2 + \frac{K_0^2(4 + C^2R^2)}{2BR^2}\right]\delta\alpha(t), \end{aligned} \tag{41}$$

where $\delta\alpha(t) = -\delta K(t)/K_0^2$. Using Eqs. (34) and (37) , we can also express the absorbed energy in terms B and C as

$$\langle \hat{H}_q \rangle - \langle \hat{H}_q \rangle_{t=0} = \frac{K_0(4 + C^2R^2)}{16B} + \frac{q^2R^2B}{4K_0} - \frac{qR}{2}. \tag{42}$$

As expected, for $R = 1$ ($T_0 = 0$), Eqs. (41) and (42) above are respectively equivalent to Eqs. (28) and (3) through a change of variables: $B = 1/q\Re z_q$ and $C = -2\Im z_q/\Re z_q$. Applying the Fokker-Planck approach of Sec. VI to Eqs. (41) and (42), we obtain $\overline{\langle \epsilon_q \rangle} = \frac{qR}{2}\left(\exp\left[2k_0^2W^2q^2t\right] - 1\right)$, and find that the noise-averaged absorbed energy depends on the initial temperature only through the prefactor $R = \coth\left(\frac{q}{2k_BT_0}\right)$.

VIII. NOISE-AVERAGED CORRELATION FUNCTIONS

In this section, working in the limit of small δz_q, we consider the effect of noise on the following noise-averaged correlation functions:

$$\overline{\mathcal{C}_1(x-x')} \equiv \overline{\langle \Phi(x)\Phi(x') \rangle}, \qquad \overline{\mathcal{C}_2(x-x')} \equiv \overline{\langle \partial_x\Phi(x)\,\partial_{x'}\Phi(x') \rangle}, \qquad \overline{\mathcal{C}_3(x-x')} \equiv \overline{\langle e^{i\nu\Phi(x)}\,e^{-i\nu\Phi(x')} \rangle}. \tag{43}$$

Notice that even though, in the absence of noise (or for each realization of noise), the three correlation functions above are related by simple differentiation or exponentiation, i.e., $\overline{\mathcal{C}_2(x-x')} = \overline{\partial_x\partial_{x'}\,\mathcal{C}_1(x-x')}$ and $\overline{\mathcal{C}_3(x-x')} = \overline{e^{\frac{\nu^2}{2}\mathcal{C}_1(x-x')}}$, the noise averaging must be done at the last step, and such simple relations do not necessarily hold for noise-averaged quantities.

Let us start with $\mathcal{C}_1(x) = \frac{1}{L}\sum_{q>0}\frac{\cos(qx)}{q\Re z_q}$ [see Eq. (6)]. Bearing in mind that the noise average of an odd power of δz_q vanishes, to leading order in δz_q, we can write

$$\overline{\mathcal{C}_1(x)} \approx \frac{1}{L}\sum_{q>0} K_0\frac{\cos(qx)}{q}\left(1 + K_0^2\,\overline{(\Re\delta z_q)^2}\right). \tag{44}$$

We can now write $\overline{(\Re\delta z_q)^2} = 4q^2W^2\int_0^t \sin^2\left[2q(t'-t)\right]dt' = \frac{W^2}{2}q\left[4qt - \sin(4qt)\right]$, which yields

$$\overline{\mathcal{C}_1(x)} \approx \frac{1}{2\pi}\left(K_0\int_\epsilon^\pi dq\frac{\cos(qx)}{q} + \frac{K_0^3W^2}{2}\int_0^\pi dq\cos(qx)\left[4qt - \sin(4qt)\right]\right), \tag{45}$$

where $\epsilon \simeq \frac{2\pi}{L}$ is a short-distance momentum cutoff. The first term in the right-hand side of the above equation is the correlation function $\mathcal{C}_1^{(0)}(x)$ in the ground state [see Eq. (7)]. We can then perform the integral in the second term and write

$$\overline{\mathcal{C}_1(x)} - \mathcal{C}_1^{(0)}(x) = \frac{K_0^3 W^2}{4\pi}\left(\frac{8t-(4t+x)\cos[\pi(4t-x)]-(4t-x)\cos[\pi(4t+x)]}{2x^2-32t^2} + \frac{4t[\cos(\pi x)+\pi x\sin(\pi x)-1]}{x^2}\right). \tag{46}$$

The right-hand side of the above equation simplifies by neglecting fast oscillatory terms for $x \gg 1$, and using $x \gg t$ (small δz_q requires finite t):

$$\overline{\mathcal{C}_1(x)} - \mathcal{C}_1^{(0)}(x) \approx \frac{K_0^3 W^2}{4\pi}\left(\frac{8t}{2x^2-32t^2} - \frac{4t}{x^2}\right) \approx \frac{16}{\pi} K_0^3 W^2 \frac{t^3}{x^4}. \tag{47}$$

Notice that we could obtain the same result, without an exact computation, by noting that, in the asymptotic limit of large x, the main contribution comes from small momenta. We can then expand $4qt - \sin(4qt) \simeq (4qt)^3/3$ in Eq. (45) for small q, and compute a much simpler integral, which directly leads to Eq. (47).

Similarly, we can write

$$\overline{\mathcal{C}_2(x)} \approx \frac{1}{L}\sum_{q>0} K_0\, q \cos(qx)\left(1 + K_0^2\, \overline{(\Re\delta z_q)^2}\right), \tag{48}$$

which leads to

$$\overline{\mathcal{C}_2(x)} \approx \frac{1}{2\pi}\left(K_0 \int_0^\pi dq\, q\cos(qx) + \frac{K_0^3 W^2}{2}\int_0^\pi dq\, q^2\, \cos(qx)\,[4qt - \sin(4qt)]\right). \tag{49}$$

The first integral above gives $\mathcal{C}_2^{(0)}(x) = \frac{K_0}{2\pi}\left[-1+\cos(\pi x)+\pi x\sin(\pi x)\right]/x^2$, which after neglecting the fast oscillatory terms reduced to the usual $-\frac{K_0}{2\pi}\frac{1}{x^2}$ scaling of the current-current correlation functions. By performing the second integral, and similarly neglecting the fast oscillatory terms, we can write the noise-induced correction as

$$\overline{\mathcal{C}_2(x)} - \mathcal{C}_2^{(0)}(x) \approx \frac{K_0^3 W^2}{4\pi}\left(24\frac{t}{x^4} + \frac{1}{(x+4t)^3} - \frac{1}{(x-4t)^3}\right) \approx -\frac{320}{\pi} K_0^3 W^2 \frac{t^3}{x^6}. \tag{50}$$

The calculation of $\overline{\mathcal{C}_3(x)}$ is also possible by exactly the same method. The first step is to expand the correlation function to second order in δz_q. We begin by writing

$$\overline{\mathcal{C}_3(x)} \approx \overline{e^{\frac{\nu^2}{2L}\sum_{q>0} K_0 \frac{\cos(qx)}{q}\left(1 - K_0\,(\Re\delta z_q) + K_0^2\,(\Re\delta z_q)^2\right)}}, \tag{51}$$

which gives

$$\begin{aligned}\overline{\mathcal{C}_3(x)} &\approx \mathcal{C}_3^{(0)}(x)\overline{\prod_{q>0}\left[1 + \frac{\nu^2}{2L}K_0^2\frac{\cos(qx)}{q}\left(-\Re\delta z_q + K_0\,(\Re\delta z_q)^2\right) + \frac{\nu^4}{8L^2}K_0^4\frac{\cos^2(qx)}{q^2}\,(\Re\delta z_q)^2\right]} \\ &\approx \mathcal{C}_3^{(0)}(x)\left\{1 + \frac{\nu^2}{2L}K_0^3\sum_{q>0}\frac{\cos(qx)}{q}\,\overline{(\Re\delta z_q)^2} + \frac{\nu^4}{4L^2}K_0^4\sum_{q_1,q_2>0}\frac{\cos(q_1 x)\cos(q_2 x)}{q_1 q_2}\overline{(\Re\delta z_{q_1})(\Re\delta z_{q_2})}\right\}.\end{aligned} \tag{52}$$

Notice that to go from the first to the second line, we have neglected a term of the form $\frac{1}{L^2}\sum_{q>0}$ (which scales as $\frac{1}{L}$). We can also write

$$\overline{(\Re\delta z_{q_1})(\Re\delta z_{q_2})} = 4q_1 q_2 W^2 \int_0^t \sin\left[2q_1(t'-t)\right]\sin\left[2q_2(t'-t)\right]dt' = q_1 q_2 W^2\left(\frac{\sin[2(q_1-q_2)t]}{q_1-q_2} - \frac{\sin[2(q_1+q_2)t]}{q_1+q_2}\right). \tag{53}$$

For large x, we can expand the above expression in small momenta, and obtain $\overline{(\Re\delta z_{q_1})(\Re\delta z_{q_2})} \simeq \frac{32}{3}W^2 q_1^2 q_2^2 t^3$, which upon performing the integrals gives

$$\overline{\mathcal{C}_3(x)} - \mathcal{C}_3^{(0)}(x) \sim t^3/x^{4+\nu^2/2},$$

with $\mathcal{C}_3^{(0)}(x) = e^{\frac{\nu^2}{2L}\sum_{q>0} K_0 \frac{\cos(qx)}{q}} \propto x^{-\frac{\nu^2}{2}}$.